

## SMAD3

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC ICC IP

**Recommended Dilution:**WB 1:500 - 1:1000 IHC 1:50 - 1:200 ICC 1:50 - 1:200 IP 1:20 - 1:100

**Calculated MW:**58kDa

**Observed MW:**Refer to Figures

**Immunogen:**

A synthetic peptide of human SMAD3

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

r

**Synonym:**

HSPC193; HsT17436; JV15-2; LDS1C; LDS3; MADH3;

**Catalog #:**A0427

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**4088

**Isotype:**IgG

**Swiss Prot:**P84022

**Purity:**Affinity purification

For research use only.

**Background:**

Smad proteins belong to a group of intracellular signal transducers and downstream effectors of TGF-/BMP Signaling. The family consists of nine members that are highly conserved in the N- and C-terminal regions. Smad3 is a key component in intracellular signaling of transforming growth factor beta (TGF-), an inhibitor for tumor cell proliferation. Smad3 is activated by activin/TGF- receptors, which form a heteromeric complex with Smad4, translocating to the nucleus and regulating the expression of TGF- target genes. Interaction of Smad3 with Akt, can initiate TGF- induced apoptosis and cell cycle arrest. Smad3 expression may be critical in tumor suppression in early stages of gastric carcinogenesis.

*To place an order, please [Click HERE](#).*