

Smad2

Reactivity: Human Mouse

Tested applications: WB ICC

Recommended Dilution: WB 1:500 - 1:2000 ICC 1:50 - 1:200

Calculated MW: 58kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant protein of human Smad2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

1 mg/ml

Synonym:

SMAD2; JV18 ;JV18-1 ; MADH2 ; MADR2 ;MGC22139 ;MGC34440 ; hMAD-2 ;hSMAD2;

Catalog #: A0441

Antibody Type:

Monoclonal Antibody

Species: Mouse

Gene ID: 4087

Isotype: IgG

Swiss Prot: Q15796

Purity: Affinity purification

For research use only.

Background:

Members of the Smad family of signal transduction molecules are components of a critical intracellular pathway that transmit TGF- signals from the cell surface into the nucleus. Three distinct classes of Smads have been defined: the receptor-regulated Smads (R-Smads), which include Smad1, 2, 3, 5, and 8; the common-mediator Smad (co-Smad), Smad4; and the antagonistic or inhibitory Smads (I-Smads), Smad6 and 7 (1-5). Activated type I receptors associate with specific R-Smads and phosphorylate them on a conserved carboxy-terminal SSXS motif. The phosphorylated R-Smad dissociates from the receptor and forms a heteromeric complex with the co-Smad (Smad4), allowing translocation of the complex to the nucleus. Once in the nucleus, Smads can target a variety of DNA binding proteins to regulate transcriptional responses (6-8).

To place an order, please [Click HERE](#).