

## CDC25C

**Reactivity:**Human Mouse

**Tested applications:**WB IF

**Recommended Dilution:**WB 1:500 - 1:2000 IF 1:50 - 1:200

**Calculated MW:**53kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human CDC25C

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

CDC25; PPP1R60;

**Catalog #:**A11328

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**995

**Isotype:**IgG

**Swiss Prot:**P30307

**Purity:**Affinity purification

For research use only.

**Background:**

Cdc25 is a protein phosphatase responsible for dephosphorylating and activating cdc2, a crucial step in regulating the entry of all eukaryotic cells into mitosis (1). cdc25C is constitutively phosphorylated at Ser216 throughout interphase by c-TAK1, while phosphorylation at this site is DNA damage-dependent at the G2/M checkpoint (2). When phosphorylated at Ser216, cdc25C binds to members of the 14-3-3 family of proteins, sequestering cdc25C in the cytoplasm and thereby preventing premature mitosis (3). The checkpoint kinases Chk1 and Chk2 phosphorylate cdc25C at Ser216 in response to DNA damage (4,5).

*To place an order, please [Click HERE](#).*