

USP22

Reactivity:Human

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:1000 IHC 1:50 - 1:100

Calculated MW:59kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human USP22

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

f

Synonym:

USP3L;

Catalog #:A6173

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:23326

Isotype:IgG

Swiss Prot:Q9UPT9

Purity:Affinity purification

For research use only.

Background:

Modification of target proteins by ubiquitin participates in a wide array of biological functions. Proteins destined for degradation or processing via the 26 S proteasome are coupled to multiple copies of ubiquitin. However, attachment of ubiquitin or ubiquitin-related molecules may also result in changes in subcellular distribution or modification of protein activity. An additional level of ubiquitin regulation, deubiquitination, is catalyzed by proteases called deubiquitinating enzymes, which fall into four distinct families. Ubiquitin C-terminal hydrolases, ubiquitin-specific processing proteases (USPs), 1 OTU-domain ubiquitin-aldehyde-binding proteins, and Jab1/Pad1/MPN-domain-containing metallo-enzymes. Among these four families, USPs represent the most widespread and represented deubiquitinating enzymes across evolution. USPs tend to release ubiquitin from a conjugated protein. They display similar catalytic domains containing conserved Cys and His boxes but divergent N-terminal and occasionally C-terminal extensions, which are thought to function in substrate recognition, subcellular localization, and protein-protein interactions.

To place an order, please [Click HERE](#).