

IFN g Human

Description: Interferon-gamma Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 144 amino acids and having a molecular mass of 16879 Dalton. The IFN-gamma is purified by proprietary chromatographic techniques.

Synonyms: Immune Interferon, type II interferon, T cell interferon, MAF, IFNG, IFG, IFI, IFN-gamma.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: MQDPYVKEAE NLKKYFNAGH SDVADNGTLF LGILKNWKEE
SDRKIMQSQI VSFYFKLFKN FKDDQSIQKS VETIKEDMNV KFFNSNKKKR DDFEKLTNYS
VTDLNVQRKA IHELIQVMAE LSPAAGTGKR KRSQMLFQGR RASQ.

Purity: Greater than 98.0% as determined by (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

Formulation:

The protein was lyophilized after extensive dialysis from a concentrated (1mg/ml) solution in 10mM sodium Phosphate buffer pH=7.4

Stability:

Lyophilized Interferon gamma although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IFN-gamma should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized Interferon-gamma in sterile 18M-cm H₂O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

Introduction:

IFN-gamma produced by lymphocytes activated by specific antigens or mitogens. IFN-gamma, in addition to having antiviral activity, has important immunoregulatory functions, it is a potent activator of macrophages, and has antiproliferative effects on transformed cells and it can potentiate the antiviral and antitumor effects of the type I interferons.

Biological Activity:

The specific activity as determined in a viral resistance assay is ≥ 0.05 ng/ml, corresponding to a specific activity of $2.0 \times 10,000,000$ IU/mg.

References:

Title: Candida albicans abrogates the expression of interferon--inducible protein-10 in human

www.neobiolab.com

info@neobiolab.com

888.754.5670, +1 617.500.7103 United States

0800.088.5164, +44 020.8123.1558 United Kingdom

keratinocytes.Publication:Article first published online: 18 JUL 2008DOI:

10.1111/j.1574-695X.2008.00457.xLink: <http://onlinelibrary.wiley.com/doi/10.1111/j.1574-695X.2008.00457.x/full>

8.00457.x/full



Catalog #:CYPs-213

To place an order, please [Click HERE](#).

For research use only.

© 2015 NeoScientific



neobiolab | Term | Quote/Order | Services | Products | Support | Corporate | Contact Us

Copyrights © 2013 NEO Group 245 First Street, 18th Floor, Cambridge MA 02142 888.754.5670 (toll-free)



Request a
CATALOG
[CLICK HERE](#)