

GH Bovine

Description: Growth Hormone Bovine Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 191 amino acids and having a molecular mass of 21.8 kDa. The GH Bovine Recombinant is purified by proprietary chromatographic techniques.

Synonyms: BGH, BST, rBGH, rBST, Bovine Somatotropin, Bovine GH, Growth hormone, GH1.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: afpamsls gl fanavraqh lhqlaadtfk efertyiepg qrysiqntqv afcfsetmpa
ptgkneaqqk sdlellrlsl lliqswlgpl qflsrvtfs lvfgtsdrvy eklkdleegi lalmreledg tprrgqilkq tydkfdtnmr
sddallknvg llscfrkdlh ktetylrvmk crfgeasca f.

Purity: Greater than 98.0% as determined by: (a) Analysis by SEC-HPLC. (b) Analysis by SDS-PAGE.

Formulation:

The Bovine Growth Hormone protein was lyophilized from a concentrated (1mg/ml) solution with 0.0045mM NaHCO₃ adjusted to pH-8.

Stability:

Lyophilized Bovine Growth Hormone although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution GH Bovine should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized Bovine Growth Hormone in sterile 18M-cm H₂O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

Introduction:

GH is a member of the somatotropin/prolactin family of hormones which play an important role in growth control. The gene, along with four other related genes, is located at the growth hormone locus on chromosome 17 where they are interspersed in the same transcriptional orientation; an arrangement which is thought to have evolved by a series of gene duplications. The five genes share a remarkably high degree of sequence identity. Alternative splicing generates additional isoforms of each of the five growth hormones, leading to further diversity and potential for specialization. This particular family member is expressed in the pituitary but not in placental tissue as is the case for the other four genes in the growth hormone locus. Mutations in or deletions of the gene lead to growth hormone deficiency and short stature.

Biological Activity:

The activity as determined by the PDFP13B9 cells stably transfected with rabbit GH receptors.

www.neobiolab.com

info@neobiolab.com

888.754.5670, +1 617.500.7103 United States

0800.088.5164, +44 020.8123.1558 United Kingdom

Bovine GH is also capable of forming a 1:2 complex with the recombinant ovine growth hormone receptor extracellular domain (ECD).



To place an order, please [Click HERE](#).

Catalog #:CYP5-643

For research use only.

© 2015 NeoScientific



neobiolab | [Term](#) | [Quote/Order](#) | [Services](#) | [Products](#) | [Support](#) | [Corporate](#) | [Contact Us](#)

Copyrights © 2013 NEO Group 245 First Street, 18th Floor, Cambridge MA 02142 888.754.5670 (toll-free)



Request a
CATALOG
[CLICK HERE](#)