

IL 28B Mouse

Description:IL-28B Mouse Recombinant produced in E.coli is a single, non-glycosylated polypeptide chain containing 174 amino acids and having a molecular mass of 19.6kDa. The IL28B is purified by proprietary chromatographic techniques.

Catalog #:CYP5-739

Synonyms:Interferon lambda-3, IFN-lambda-3, Interleukin-28B, IL-28B, Ifnl3, Il28, Il28b, IFL-1, INF-alpha, INF-lambda.

For research use only.

Source:Escherichia Coli.

Physical Appearance:Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence:DPVPRATRLP VEAKDCHIAQ FKSLSPKELQ AFKKAKGAIE
KRLLEKDMRC SSSLISRAWD LKQLVQERP KALQAEVALT LKVVENINDS ALTTILGQPL
HTLSHIHSQL QTCTQLQATA EPKPPSRRLS RWLHRLQEAQ SKETPGCLEL SVTSNLFQLL
LRDLKCVASG DQCV.

Purity:Greater than 97.0% as determined by: (a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Formulation:

IL28B protein was lyophilized from a 0.2

Stability:

Lyophilized IL-28B although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL28B should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolabs products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized IL-28B in sterile 18M-cm H₂O not less than 100

Introduction:

IL-28B is a cytokine remotely related to type I interferons and the IL-10 family. Interleukin-28B, interleukin 28A (IL28A), and interleukin 29 (IL29) are 3 closely related cytokine genes which form a cytokine gene cluster on a chromosomal region mapped to 19q13. Expression of the cytokines encoded by these 3 genes can be induced by viral infection. All 3 cytokines interact with a heterodimeric class II cytokine receptor which consists of interleukin 10 receptor, beta (IL10RB) and interleukin 28 receptor, alpha (IL28RA).

Biological Activity:

The ED50 as determined by an anti-viral assay using human HepG2 cells infected with encephalomyocarditis is less than 30 ng/ml, corresponding to a specific activity of > 3.3

To place an order, please [Click HERE](#).