

SIRT3 Human

Description: SIRT3 produced in E.Coli is a single, non-glycosylated polypeptide chain containing 303 amino acids (118-399a.a.) and having a molecular mass of 33.5kDa. SIRT3 is fused to a 21 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #: PRPS-469

For research use only.

Synonyms: Sirtuin 3, SIR2-like protein 3, Sir2-like 3, NAD-dependent deacetylase sirtuin-3, mitochondrial.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered clear solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MSDKGKLSLQ DVAELIRARA
CQRVVMVGA GISTPSGIPD FRSPGSGLYS NLQYDLPYP EAIFELPFFF HNPKEPFTLA
KELYPGNYKP NVTHYFLRL HDKGLLLRLY TQNDGLERV SGIPASKLVE AHGTFASATC
TVCQRPFPGE DIRADVMDR VPRCPVCTGV VKPDIVFFGE PLPQRFLHV VDFPMADLLL
ILGTSLEVEP FA

Purity: Greater than 95% as determined by SDS-PAGE.

Formulation:

The SIRT3 protein solution (1mg/1ml) is formulated in 20mM Tris-HCl buffer (pH8.0) 1mM DTT, 0.1M NaCl, 10% glycerol.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

SIRT3 is a member of a family of proteins called Sirtuin. Proteins of the Sirtuin family are characterized by a Sirtuin core domain and grouped into four classes. The roles of human Sirtuins are numerous and are important in aging processes, stress resistance and metabolic regulation. SIRT3 shows NAD⁺-dependent deacetylase activity in the mitochondria. Over-expression of SIRT3 causes increased levels of the mitochondrial uncoupling protein 1. Also, in certain breast cancers the levels of SIRT3 protein are found to be extremely high.

To place an order, please [Click HERE](#).